

Household Energy Consumption, Emissions, Pollution, and Health Impacts in India

STATE Dadra & Nagar Haveli

(state and district as of census-India, 2011)

DISTRICT Dadra_Nagar_Haveli

Household energy consumption (HEC) emissions were calculated in four classes - cooking (CK), lighting (LG), space heating (SH), and water heating (WH). Bottom-up emissions for the four classes are available @ 0.25 degree spatial resolution, and further aggregated to district and state level. A sub-classification is available by fuel - biomass, coal, kerosene, liquified petroleum gas (LPG), and others.



%Households Primary Cooking Fuel

gas+elec

others

40.6%

59.4%

Estimated district annual HEC emissions

| | |
|----------------------|---------------|
| Paticulates (2.5µm) | 160 tons |
| Sulfur dioxide | 20 tons |
| Nitrogen oxides | 5 tons |
| Carbon monoxide | 2,800 tons |
| Hydrocarbons | 300 tons |
| Black carbon (BC) | 40 tons |
| Organic carbon | 80 tons |
| Carbon dioxide (CO2) | 0.01 mil tons |


Estimated PM_{2.5} emissions @ 0.25 degree resolution




Modeled share of HEC emissions to ambient PM_{2.5}





DISTRICT EMISSIONS





NATIONAL EMISSIONS



CO₂



BC



% contribution of HEC emissions to modeled ambient PM_{2.5} concentrations

(concentrations were conducted using the WRF-CAMx models)

| | |
|----------|-------|
| National | 29.6% |
| District | 19.3% |

The health impacts of outdoor air pollution as ischemic heart diseases (which can lead to heart attacks), cerebrovascular disease (which can lead to strokes), chronic obstructive pulmonary diseases, lower respiratory infections, and cancers (in trachea, lungs, and bronchitis) were estimated using the age-dependent relative risk functions detailed in the Global Burden of Disease study and dispersion modeling results from this study. The final calculations were conducted at the district level using the population distribution by age presented in Census-India.

Estimated premature mortality of outdoor air pollution per year - apportioned to HEC emissions

| | |
|----------|------------------|
| National | 84,000 - 115,000 |
| District | 6 - 7 |

Emission and dispersion modeling results, pollution animations, and summary sheets by district and state are hosted @ <http://www.urbanemissions.info>
Send your comments and questions to sim-air@urbanemissions.info