

# Household Energy Consumption, Emissions, Pollution, and Health Impacts in India


STATE **Haryana**

(state and district as of census-India, 2011)

DISTRICT **Ambala**

Household energy consumption (HEC) emissions were calculated in four classes - cooking (CK), lighting (LG), space heating (SH), and water heating (WH). Bottom-up emissions for the four classes are available @ 0.25 degree spatial resolution, and further aggregated to district and state level. A sub-classification is available by fuel - biomass, coal, kerosene, liquified petroleum gas (LPG), and others.

### Emissions and Dispersion Modeling Schematics



### %Households Primary Cooking Fuel

gas+elec

others


59.2%

40.8%


### Estimated district annual HEC emissions

Paticulates (2.5µm)	2,440 tons
Sulfur dioxide	290 tons
Nitrogen oxides	55 tons
Carbon monoxide	24,000 tons
Hydrocarbons	6,040 tons
Black carbon (BC)	380 tons
Organic carbon	900 tons
Carbon dioxide (CO <sub>2</sub> )	0.19 mil tons


### Estimated PM<sub>2.5</sub> emissions @ 0.25 degree resolution








### Modeled share of HEC emissions to ambient PM<sub>2.5</sub>



### DISTRICT EMISSIONS



### NATIONAL EMISSIONS



### % contribution of HEC emissions to modeled ambient PM<sub>2.5</sub> concentrations

(concentrations were conducted using the WRF-CAMx models)

National	29.6%
District	28.7%

The health impacts of outdoor air pollution as ischemic heart diseases (which can lead to heart attacks), cerebrovascular disease (which can lead to strokes), chronic obstructive pulmonary diseases, lower respiratory infections, and cancers (in trachea, lungs, and bronchitis) were estimated using the age-dependent relative risk functions detailed in the Global Burden of Disease study and dispersion modeling results from this study. The final calculations were conducted at the district level using the population distribution by age presented in Census-India.

### Estimated premature mortality of outdoor air pollution per year - apportioned to HEC emissions

National	84,000 - 115,000
District	49 - 56

Emission and dispersion modeling results, pollution animations, and summary sheets by district and state are hosted @ <http://www.urbanemissions.info>  
Send your comments and questions to [sim-air@urbanemissions.info](mailto:sim-air@urbanemissions.info)